Ap. J. Lett., in press
Preprint typeset using LaTeX style emulateapj v. 04/03/99

arXiv:astro-ph/9907023v2 14 Oct 1999

THERMAL STABILITY OF COLD CLOUDS IN GALAXY HALOS

Mark Wardle & Mark Walker
Special Research Centre for Theoretical Astrophysics,
School of Physics, University of Sydney, NSW 2006, Australia
Ap. J. Lett., in press

ABSTRACT

We consider the thermal properties of cold, dense clouds of molecular hydrogen and atomic helium. For cloud masses below $10^{-1.7}\mathrm{M}_\odot$, the internal pressure is sufficient to permit the existence of particles of solid or liquid hydrogen at temperatures above the microwave background temperature. Optically thin thermal continuum emission by these particles can balance cosmic-ray heating of the cloud, leading to equilibria which are thermally *stable* even though the heating rate is independent of cloud temperature. For the Galaxy, the known heating rate in the disk sets a minimum mass of order $10^{-6}\mathrm{M}_\odot$ necessary for survival. Clouds of this type may in principle comprise most of the dark matter in the Galactic halo. However, we caution that the equilibria do not exist at redshifts $z \gtrsim 1$ when the temperature of the microwave background was substantially larger than its current value; the formation and survival of such clouds to the present epoch therefore remain open questions.

Subject headings: ISM: clouds — galaxies: halos — dark matter

1. INTRODUCTION

Walker & Wardle (1998) showed that a population of neutral, AU-sized clouds in the Galactic halo could be responsible for the "Extreme Scattering Events" (ESEs) observed in the radio flux towards several quasars (Fiedler et al. 1987, 1994). In this model the cloud surfaces are exposed to UV radiation from hot stars in the Galactic disk, producing a photo-ionised wind. When one of these clouds crosses the line of sight to a compact radio source, the flux varies as a result of refraction by the ionised gas (cf. Henriksen & Widrow 1995). This model explains the observed flux variations quite naturally; but if the clouds are self-gravitating, then the ESE event rate implies that the cloud population comprises a significant fraction of the Galaxy's mass.

This halo cloud population cannot contain much dust mixed with the gas as this would lead to optical extinction events of distant stars: either the clouds have extremely low metallicity, or any dust grains have sedimented to the cloud centre. Given this, several factors make the clouds difficult to detect (Pfenniger, Combes & Martinet 1994): cold molecular hydrogen is, by and large, invisible; the clouds are small; they are transparent in most regions of the electromagnetic spectrum; and they cover a small fraction of the sky. The clouds are not sufficiently compact to cause gravitational lensing towards the LMC, although Draine (1998) has shown that there is substantial optical refraction by the neutral gas, so that microlensing experiments (Paczyński 1996) already place useful constraints on the properties of low-mass halo clouds.

Given that this hypothesised cloud population does not violate observational constraints, the primary issues that need to be addressed are theoretical: (i) how and when did these clouds form? and (ii) how do they resist gravitational collapse? The second of these is addressed in this *Letter*.

We begin by writing down equations describing a simple "one-zone" model of a cloud, characterised by a single temperature and pressure (§2), and show that particles of solid H_2 may exist in the clouds (cf. Pfenniger & Combes 1994). At temperatures above the microwave background temperature, these particles cool the cloud by thermal continuum radiation, admitting equilibria in which this cooling balances heating by cosmic rays. In §3, we show that (for optically thin emission) these equilibria are thermally stable: if the cloud contracts the coolant is destroyed by the increase in temperature, and the power deposited by cosmic rays causes the cloud to expand and the temperature to return to its original value. We conclude that, within the context of our one-zone model, the viable mass range for Galactic clouds is 10^{-6}–$10^{-1.7}\mathrm{M}_\odot$.

2. CLOUD MODEL

Virial equilibrium implies that for a self-gravitating cloud characterised by mass M, temperature T, and radius R,

$$R \approx GM\mu/kT \tag{1}$$

where μ is the mean molecular weight. The pressure in the cloud can be related to the temperature upon noting that $P \sim GM^2/R^4$, yielding

$$P = \frac{q}{G^3 M^2}\left(\frac{kT}{\mu}\right)^4 , \tag{2}$$

where q depends on the cloud's structure. For polytropes q rises monotonically from 9.2 to 40 as the polytropic index runs from 3/2 to 9/2, so we adopt $q = 20$.

At sufficiently high pressures a fraction x of the molecular hydrogen assumes solid (or liquid) form. Then $\mu = (1 - x + 2y)m/(1 - x + y)$, where m is the mass of an H_2 molecule, and $y \approx 1/6$ is the abundance ratio He:H_2 by number. Neglecting the temperature difference between the phases, in equilibrium the partial pressure of H_2 equals the saturated vapour pressure, i.e.

$$\frac{1-x}{1-x+y}P = \left(\frac{2\pi m}{h^2}\right)^{3/2} (kT)^{5/2} e^{-T_v/T} , \tag{3}$$

(valid for $0 < x < 1$), where kT_v is the heat of vapourisation for H_2 (Phinney 1985). With $T_v = 91.5\,\mathrm{K}$, the vapour pressure given by the RHS of eq. (3) is within 20 % of the available experimental data (Souers 1986).

Hydrogen grains can cool the gas in a manner similar to dust grains in molecular clouds: the gas cools via collisions with



FIG. 1.— *Top:* cooling by rotational transitions of H_2, HD and LiH in the gas phase, and thermal continuum emission from particles of solid H_2 for a $10^{-3}M_{\odot}$ cloud in virial equilibrium. *Bottom:* solid H_2 cooling rates for cloud masses 10^{-3}, 10^{-5} and $10^{-7}M_{\odot}$. Dotted portion of the $10^{-7}M_{\odot}$ curve indicates where the coolant is liquid H_2. The dashed line in both plots indicates the cooling rate that would give a Kelvin-Helmholtz time scale of 10 Gyr.

slightly colder solid particles, which in turn cool by thermal continuum emission. To calculate the cooling by solid H_2, first consider the net power radiated by a single particle (we employ an 'escape probability' formulation of radiative transfer):

$$L_s = 4\sigma \left(\frac{C(T)T^4}{1+\tau} - \frac{C(T_b)T_b^4}{1+\tau_b} \right) , \tag{4}$$

where $C(T)$ is the Planck-mean absorption cross-section, T_b is the cosmic microwave background temperature, and τ and $\tau_b = \tau C(T_b)/C(T)$ are the Planck-mean optical depth of the cloud to thermal radiation characterized by T and T_b respectively. Assuming that the particle size is $\ll \lambda$ ($\sim 0.1\,\mathrm{cm}$ at the temperatures of relevance here), we may write $C(T) = C_m(T)m_s$, m_s being the particle mass, and for spherical grains we have (Draine & Lee 1984)

$$C_m(T) \approx \frac{15(4\pi)^3}{28\rho_s} \frac{\lambda_2}{(\varepsilon_1+2)^2} \left(\frac{kT}{hc} \right)^2 , \tag{5}$$

where $\rho_s = 0.087\,\mathrm{g\,cm^{-3}}$ is the density of solid H_2 (Souers 1986),the complex dielectric function of the solid is $\varepsilon_1 + i\varepsilon_2$, and we have assumed that $\varepsilon_2 = \lambda_2/\lambda$ as expected at low frequencies. The net cooling rate per unit mass of cloud material (gas and solid) is then

$$\Lambda = \frac{4\pi R^2\sigma}{M} \left(\frac{\tau T^4}{1+\tau} - \frac{\tau_b T_b^4}{1+\tau_b} \right) , \tag{6}$$

where $\tau_b = \tau\,(T_b/T)^2$, and

$$\tau = C_m \frac{x}{1+2y} \frac{M}{\pi R^2} . \tag{7}$$

To evaluate Λ, we require optical constants for solid H_2 in the microwave. The particles are expected to be almost pure para-hydrogen as an ortho-para mixture of the solid relaxes to para ($J = 0$) form in a few days (Souers 1986). The low frequency value of ε_1 for para-hydrogen has been measured (Souers 1986) as $\varepsilon_1 \simeq 1.25$; the low frequency limit of ε_2 is less certain. Jochemsen et al. (1978) measured the extinction coefficient of a single crystal of solid para-hydrogen in the region of interest ($\lambda \sim 0.1$ cm), but could not determine whether this continuum extinction was due to absorption or scattering within the crystal. Because these measurements do not conform to the anticipated low-frequency behaviour ($\propto 1/\lambda$), and absorption bands are not expected below the S(0) line, it is likely that the absorption of pure crystalline H_2 is much smaller than the measured extinction, and we can only infer a limit: $\varepsilon_2 \lesssim 1.8 \times 10^{-3}$. However, the low-frequency absorption of solid H_2 grains could be strongly enhanced by impurity species and lattice defects. For the purposes of this paper we adopt $\varepsilon_2 = \lambda_2/\lambda$ and $\lambda_2 = 10^{-4}\,\mathrm{cm}$. Within the confines of the model, this assumption represents one of our main areas of uncertainty.

For a given cloud mass, the fraction of H_2 in the solid phase can be determined from T using eqs (2) and (3). This allows the cooling by solid particles to be calculated as a function of T. The upper panel of Fig 1 illustrates this for a cloud of mass $10^{-3}\mathrm{M}_\odot$. For comparison, the cooling contributed by rotational lines of gas-phase H_2, HD and LiH is also plotted; energies and A-values from Turner, Kirby-Docken & Dalgarno (1977), Abgrall, Roueff & Viala (1982), and Gianturco et al. (1996). The adopted deuterium and LiH abundances are 3×10^{-5}, and 1.2×10^{-10} respectively (Schramm & Turner 1998). We employ an escape-probability formulation of radiative transfer, in which the optically-thin cooling rates are divided by $(1+\tau)$ where τ is the optical depth at line centre (rotational transitions) or the Planck-mean (continuum emission).

There is a critical temperature T_c at which H_2 in the cloud lies on the border between the solid and gaseous phases, i.e. the partial pressure of H_2 is equal to its saturated vapour pressure. From this point x and Λ increase precipitously as T is reduced, until $\tau \sim 1$ and the cooling is then roughly black-body. Λ then increases more slowly, peaking when $x \sim 1/2$ and subsequently dropping to zero as the temperature approaches that of the microwave background. At lower temperatures the cloud is heated by the background radiation.

The solid H_2 cooling curves for cloud masses of 10^{-3}, 10^{-5} and $10^{-7}\mathrm{M}_\odot$ are compared in the lower panel of figure 1. Decreasing the cloud mass has two consequences: the critical temperature T_c decreases, whereas the maximum value of Λ decreases (because in this circumstance the emission is optically thick, and cloud surface area is proportional to M^2 at a given temperature). For cloud masses $\lesssim 10^{-7}\mathrm{M}_\odot$, T_c is above the H_2 triple point (13.8 K) and liquid droplets of H_2 form instead. This does not qualitatively affect the calculations as the density and saturated vapour pressure of the liquid are within 50% of those of the solid for $T \lesssim 20\,K$ (Souers 1986), and the optical properties are similar (in the sense that ε_2 is small and uncertain), thus in Fig. 1 we continue the cooling curve for a $10^{-7}\mathrm{M}_\odot$ cloud above the triple point as a dotted curve. Thermally stable solutions do not exist for masses $\lesssim 10^{-7.5}\mathrm{M}_\odot$, as the partial pressure of H_2 exceeds the saturated vapour pressure unless $x > 0.5$ (see §3). On the other hand, for $M \gtrsim 10^{-1.7}\mathrm{M}_\odot$ T_c is below the CMB temperature and the solid phase warms the cloud rather than cooling it.

3. THERMAL EQUILIBRIUM

In thermal equilibrium, the cloud temperature is set by the balance between cooling and heating. We assume that clouds in the Galactic halo are heated primarily by cosmic rays. The local interstellar cosmic-ray ionisation rate in the Galactic disc, $\sim 3 \times 10^{-17}\,\mathrm{s^{-1}\,H^{-1}}$ (Webber 1998), implies a heating rate $\Gamma \sim 3 \times 10^{-4}\mathrm{erg\,g^{-1}\,s^{-1}}$ (Cravens & Dalgarno 1978). The cosmic-ray heating in the halo is uncertain but should be somewhat lower, say $\sim 10^{-5}\mathrm{erg\,g^{-1}\,s^{-1}}$. In Fig. 2, we show the cooling rates for clouds of masses 10^{-7}–$10^{-1.7}\mathrm{M}_\odot$. The solid curves are contours of constant optical depth; the dashed curve shows the optically-thick limit and represents the maximum cooling rate for each cloud mass. It appears that solid hydrogen can provide the necessary cooling for planetary-mass gas clouds at the cosmic-ray heating rates expected in the Galactic disk and halo.

The upper panel of Fig. 1 shows that there are typically three equilibrium temperatures available for cloud masses between $10^{-7.5}$ and $10^{-1.7}\mathrm{M}_\odot$ at the expected heating rates: solid H_2 provides one barely above T_b and one a few degrees higher; the gas-phase coolants provide an equilibrium above 30 K. We now show that thermal stability requires Λ to be a decreasing function of T, and therefore only the second of these three equilibria is stable.

In virial equilibrium the total energy per unit mass is approximately $-\frac{3}{2}kT/\mu$ (the internal excitation of the gas is negligible at the low temperatures of interest here), so the thermal evolu-



FIG. 2.— Solid H$_2$ cooling rate versus cloud mass for different optical depths. The curves, labelled by τ, are truncated at high masses when the solutions become thermally unstable. The optically thick limit, corresponding to black-body emission, is labelled "∞". The dashed curve delineates where half of the hydrogen is in solid form, an approximate upper limit for thermal stability.

tion of the cloud is determined by

$$\frac{3k}{2\mu}\frac{\mathrm{d}T}{\mathrm{d}t} = \Lambda - \Gamma. \qquad (8)$$

In the absence of heating the cloud contracts on the Kelvin-Helmholtz time-scale $t_{\rm KH} = \frac{3}{2}kT/(\mu\Lambda)$. Note that this time-scale can be a substantial fraction of the Hubble time for temperatures of a few Kelvin: the dashed curves in figure 1 show the cooling rate that yields $t_{\rm KH} = 10\,{\rm Gyr}$. In thermal equilibrium cosmic-ray heating replaces the energy radiated away by the cloud implying, for example, $t_{\rm KH} \sim 2\times10^6$ yr for $\Gamma \sim 10^{-5}\,{\rm erg\,g^{-1}\,s^{-1}}$ (at $T \sim 10$ K). This is much greater than the sound crossing time ($\sim 10^2$ yr), so the response of a cloud to dynamical perturbations is adiabatic, to a good approximation, and dynamical stability is assured. However, eq. (8) shows that perturbations to the cloud temperature grow or decay as $e^{\alpha t}$ where

$$\alpha = t_{\rm KH}^{-1}\frac{T}{\Lambda}\frac{\mathrm{d}}{\mathrm{d}T}(\Lambda-\Gamma), \qquad (9)$$

and the right-hand side of this equation is evaluated at the equilibrium temperature. Thus a cloud is thermally stable only if a decrease (increase) in cloud temperature leads to cooling outstripping (lagging) heating. For cosmic ray heating, Γ is independent of T if the column through the cloud is insufficient to cause significant attenuation of cosmic rays (changes in temperature affect the cloud's column density through the virial relationship $R \propto 1/T$). Thermal stability then requires that Λ be a decreasing function of T, and we conclude that *only the equilibrium on the high-temperature shoulder of the solid hydrogen cooling curve is stable.* In fact the column density of each cloud ($\sim 10^2\ {\rm g\,cm^{-2}}$: Walker 1999) is sufficient to stop sub-GeV cosmic-ray protons (and all electrons), leading to a dependence of Γ on T; this dependence is too weak to affect our conclusions concerning stability.

4. DISCUSSION

The suggestion that cold gas could comprise a significant fraction of the Galaxy's dark matter is not new, previous proposals include: a fractal medium in the the outer reaches of the Galactic disk (Pfenniger et al. 1994); isolated halo clouds (Gerhard & Silk 1996); and mini clusters of clouds in the halo (de Paolis et al. 1995; Gerhard & Silk 1996). However, to date there has been no compelling reason to believe that isolated, cold gas clouds – as inferred by Walker & Wardle (1998) – could support themselves for long periods against gravitational collapse. We have shown that such clouds can be stabilised by the precipitation/sublimation of particles of solid hydrogen (or by the condensation/evaporation of droplets of liquid hydrogen) if these particles dominate the radiative cooling of the cloud. The key feature which confers thermal stability is that these particles are destroyed, hence cooling becomes less efficient, as the cloud temperature increases. This feature will be present in any model where condensed hydrogen is the principal coolant, and consequently we expect that more sophisticated structural treatments will also admit stable solutions.

The masses of thermally stable clouds lie in the approximate range $10^{-7.5}$–$10^{-1.7}{\rm M}_\odot$. The lower limit is increased to $10^{-6}{\rm M}_\odot$ if subject to cosmic ray heating similar to that in the Galactic disc for an interval $\gtrsim kT/\Gamma \sim 10^5\,{\rm yr}$. As halo clouds take much longer than this to pass through the cosmic-ray disc, this limit is appropriate even for a halo cloud population. For cloud masses in the range 10^{-6}–$10^{-1.7}\ {\rm M}_\odot$ the radiative cooling simply readjusts, on the timescale $t_{\rm KH}$, to maintain equilibrium as Γ varies through the orbit ($\sim 10^8$ yr) of a cloud around the Galaxy.

The typical particle radius a is constrained by the requirement that the clouds not produce significant extinction at optical wavelengths: the geometrical optical depth of the particles, $\tau_g \approx \tau/C_m a\rho_s$ should be less than 1. Adopting $T = 5\,{\rm K}$ and $\tau = 0.01$ this translates to $a({\rm mm}) \gtrsim (10^{-4}\,{\rm cm})/\lambda_2$. Millimeter-size particles settle to the centre of the cloud in $\sim 10^4\,{\rm yr}$, this time is shortened if λ_2 is significantly less than $10^{-4}\,{\rm cm}$ and the particles are required to be larger. Settling may be counteracted by convective motions or sublimation resulting from the higher temperatures deeper within the cloud — issues that must await a more sophisticated treatment of the cloud structure.

If the hypothesised population of clouds exists, their thermal microwave emission may be detectable as a Galactic continuum background at temperatures just above the Cosmic Microwave Background; a Galactic component of this kind has in fact been isolated in the COBE FIRAS data (Reach et al. 1985). One would like to compare these data with the theory presented here, but it is difficult to predict the total microwave intensity for our model because the distribution of cosmic-ray density away from the Galactic plane is only loosely constrained (see Webber et al. 1994). A similar uncertainty afflicts the modelling of γ-ray production from baryonic material in the Galactic halo (cf. de Paolis et al. 1995; Salati et al. 1996; de Paolis et al. 1999; Kalberla, Shchekinov & Dettmar 1999). Nevertheless, microwave and γ-ray emissivities are each proportional to the local cosmic-ray flux (assuming the cosmic-ray spectrum does not vary greatly), so we can write $I_\mu \simeq I_\gamma j_\mu / j_\gamma$, for emissivities j and intensities I. At high latititudes the Galactic γ-ray background is $I_\gamma \sim 10^{-6}\ \mathrm{ph\,cm^{-2}\,s^{-1}\,sr^{-1}}$, above 1 GeV (Dixon et al. 1998); local to the Sun the corresponding (optically thin) emissivity is $1.1 \times 10^{-3}\ \mathrm{ph\ s^{-1}\,g^{-1}\,sr^{-1}}$ (Bertsch et al. 1993). Thus for a cosmic-ray heating rate (again, local to the Sun) of $\Gamma = 4\pi j_\mu \sim 3\times10^{-4}\ \mathrm{erg\,s^{-1}\ g^{-1}}$ (Cravens & Dalgarno 1978), we expect $I_\mu \sim 2\times10^{-8}\ \mathrm{erg\,cm^{-2}\,s^{-1}\ sr^{-1}}$. This is roughly 1% of that observed in the FIRAS cold component at high latitude (Reach et al. 1985), so the microwave data do not exclude the possibility of a cold cloud population heated by cosmic rays. Sciama (1999) proposed that all of the cold excess may be accounted for by cosmic-ray heating of cold clouds, but this appears to be based on an overestimate of the gamma-ray flux, and an underestimate of the high-latitude FIRAS flux.

5. CONCLUSIONS

We have demonstrated that, by virtue of the solid/gas phase transition of hydrogen, cold, planetary-mass Galactic gas clouds can be thermally stable even when they are heated at a temperature-independent rate. Our analysis applies to the present epoch, with the microwave background temperature at $T_b < 3$ K; for background temperatures $T_b \gtrsim 6$ K, our model admits no stable mass range. Consequently the longevity of the clouds at redshifts $z \gtrsim 1$ is problematic. We cannot, however, hope to address this issue until a firm theoretical basis for the formation of such clouds has been established.

We thank Sterl Phinney for making available a copy of an unpublished preprint and Bruce Draine for thoughtful comments on the manuscript. The Special Research Centre for Theoretical Astrophysics is funded by the Australian Research Council under its Special Research Centres programme.

REFERENCES

Abgrall, H., Roueff, E. & Viala, Y. 1982, A&A Supp, 50, 505
Bertsch, D. L., Dame, T. M., Fichtel, C. E., Hunter, S. D., Sreekumar, P., Stacy, J. G. & Thaddeus, P. 1993, ApJ, 416, 587
Cravens, T. E. & Dalgarno, A. 1978, ApJ, 219, 750
De Paolis, F., Ingrosso, G., Jetzer, Ph. & Roncadelli, M. 1995, A&A, 295, 567
De Paolis, F., Ingrosso, G., Jetzer, Ph. & Roncadelli, M. 1999, ApJ, 510, L103
Draine B.T. 1998, ApJ, 509, L41
Draine B.T. & Lee H.M. 1985 ApJ 285, 89
Dixon, D. D., Hartmann, D. H., Kolaczyk, E. D., Samimi, J., Diehl, R., Kanbach, G., Mayer-Hasselwander, H. & Strong, A. W. 1998, New Astronomy,3, 539
Fiedler, R. L., Dennison, B., Johnston, K. J. & Hewish, A. 1987, Nature, 326, 675
Fiedler, R., Dennison, B., Johnston, K. J., Waltman, E. B. & Simon, R. S. 1994, ApJ, 430, 581
Gerhard, O. & Silk, J. 1996, ApJ, 472, 34
Gianturco, F. A., Giorgi, P. G., Berriche, H. & Gadea, F. X. 1996, A&A Supp, 117, 377
Henriksen, R. N. & Widrow, L. M. 1995, ApJ, 441, 70
Jochemsen R., Berlinsky A.J., Verspaandonk F. & Silvera I.F. 1978, J. Low Temp. Phys 32, 185
Kalberla, P. M. W., Shchekinov, Yu. A. & Dettmar, R. J. 1999, A&A, 350, L9
Paczyński, B. 1996, ARAA, 34, 419
Pfenniger, D. & Combes, F. 1994, A&A, 285, 94
Pfenniger, D., Combes, F. & Martinet, L. 1994, A&A, 285, 79
Phinney E. S. 1985, preprint
Reach, W. T., et al. 1995, ApJ, 451, 188
Salati, P., Chardonnet, P., Luo, X. C., Silk, J. & Taillet,R. 1996, A&A, 313, 1
Schramm D.N. & Turner M.S. 1998 Rev. Mod. Phys 70, 303
Sciama D.W. 1999, MNRAS, submitted (astro-ph/9906159)
Souers, P. C. 1986, Hydrogen properties for fusion energy (Berkeley: University of California Press)
Turner, J., Kirby-Docken, K. & Dalgarno, A. 1977 ApJS 35, 281
Walker, M. 1999, MNRAS, 308, 551
Walker, M. & Wardle, M. 1998, ApJ, 498, L125
Webber, W. R. 1998, ApJ, 506, 329
Webber, W. R., Binns, W. R., Crary, D. & Westphall, M. 1994, ApJ, 429, 764